

Antimicrobial Susceptibility Profiles

- Note: The susceptibility information presented below is a summary of data gathered at ISU VDL for the time period listed. The information may be useful to understand susceptibility trends or as an aid in making clinical decisions, but may not be accurate for specific disease situations.
- In vitro antimicrobial test results do not represent therapeutic recommendations from the VDL or personnel therein. Extra/Off label usage of an antimicrobial which is limited/prohibited for certain species may result in legal action by FDA-CVM
- Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility

Avian 2008-2010	Susceptibility profile of Avian pathogens received at ISU VDL in 2008-2010					
	Data reported as: % susceptible (# isolates tested) ¹					
	E. coli	G. ana	P. mult	Salm ²	Salm B ²	Salm C1 ²
Amoxicillin	73% (168)	100% (3)	100% (1)	93% (40)	86% (22)	100% (7)
Ceftiofur	95% (168)	100% (3)	100% (1)	98% (40)	95% (22)	100% (7)
Clindamycin	0% (168)	0% (3)	0% (1)	0% (40)	0% (22)	0% (7)
Enrofloxacin	97% (168)	100% (3)	100% (1)	100% (40)	100% (22)	100% (7)
Erythromycin	0% (168)	0% (3)	0% (1)	0% (40)	0% (22)	0% (7)
Florfenicol	38% (168)	100% (3)	100% (1)	25% (40)	41% (22)	14% (7)
Gentamicin	80% (168)	100% (3)	100% (1)	83% (40)	91% (22)	71% (7)
Neomycin	85% (168)	67% (3)	100% (1)	93% (40)	95% (22)	100% (7)
Novobiocin	0% (168)	0% (3)	100% (1)	0% (40)	0% (22)	0% (7)
Oxytetracycline	36% (168)	0% (3)	100% (1)	75% (40)	59% (22)	57% (7)
Penicillin	0% (168)	0% (3)	100% (1)	0% (40)	0% (22)	0% (7)
Spectinomycin	0% (168)	0% (3)	0% (1)	0% (40)	0% (22)	0% (7)
Streptomycin	61% (168)	100% (3)	0% (1)	70% (40)	45% (22)	71% (7)
Sulfadimethoxine	47% (168)	67% (3)	0% (1)	38% (40)	32% (22)	29% (7)
Sulphathiazole	57% (168)	67% (3)	0% (1)	78% (40)	55% (22)	71% (7)
Tetracycline	36% (168)	0% (3)	100% (1)	73% (40)	59% (22)	57% (7)
Trimethoprim/Sulphamethoxazole	93% (168)	100% (3)	100% (1)	100% (40)	100% (22)	100% (7)
Tylosin (Tartrate/Base)	0% (168)	0% (3)	0% (1)	0% (40)	0% (22)	0% (7)

² See [Salmonella serotype table](#) on page 17 for most common serotypes isolated within each group

Key:

- 1 Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility
2 See *Salmonella* serotype table for most common serotypes isolated within each group
3 Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.
4 A result of <=2 ug/ml for Carbadox is a conservative indicator of bacterial inhibition by this antimicrobial agent. The result shown is based on pharmacokinetic research indicating an average Carbadox level of 4.5 mcg/ml in the small intestine of pigs fed a dose rate of 50 g/ton. (De Graff 1988).
5 Multidrug resistant isolates were found resistant to most classes of antimicrobial in the 1st round of testing. This table represents additional Disk Diffusion testing for those isolates.
- NA Not applicable
ND Not done
NI No interpretation

A equ - <i>Actinobacillus equuli</i>	H ecol - hemolytic <i>E. coli</i>	S aur - <i>Staphylococcus aureus</i>
A suis - <i>Actinobacillus suis</i>	H som - <i>Histophilus somni</i>	S beta- <i>Beta Streptococcus</i> species
Abua - <i>Acinetobacter</i> species	HPS - <i>Haemophilus parasuis</i>	S can - <i>Streptococcus canis</i>
Amy - <i>Actinomyces</i> species	K pneu - <i>Klebsiella pneumoniae</i>	S chol - <i>Salmonella choleraesuis</i>
APP - <i>Actinobacillus pleuropneumoniae</i>	M bov - <i>Moraxella bovis</i>	S dysg - <i>Streptococcus dysgalactiae</i>
B bron - <i>Bordetella bronchiseptica</i>	M haem - <i>Mannheimia haemolytica</i>	S epi- <i>Staphylococcus epidermidis</i>
B tre - <i>Bibersteinia trehalosi</i> (formerly <i>Pasteurella trehalosi</i>)	P aer - <i>Pseudomonas aeruginosa</i>	S equi - <i>Streptococcus equi</i>
Bact - <i>Bacteroides</i> group	P cab - <i>Pasteurella caballii</i>	S equus - <i>Streptococcus equisimilis</i>
C diff - <i>Clostridium difficile</i>	P mult - <i>Pasteurella multocida</i>	S pint - <i>Staph pseudintermedius</i>
C perf - <i>Clostridium perfringens</i>	Past - <i>Pasteurella</i> species	S suis - <i>Streptococcus suis</i>
Clos - <i>Clostridium</i> species	Pec - <i>Peptococcus</i> species	S ube - <i>Streptococcus uberis</i>
E coli - <i>Escherichia coli</i>	Pes - <i>Peptostreptococcus</i> species	S zoo - <i>Streptococcus zooepidemicus</i>
E fael - <i>Enterococcus faecalis</i>	Pmul A - <i>Pasteurella multocida</i> Type A	Salm sp- <i>Salmonella</i> species
E faem - <i>Enterococcus faecium</i>	Pmul D - <i>Pasteurella multocida</i> Type D	Salm B - <i>Salmonella</i> species group B
Enc - <i>Enterococcus</i> species	Prot - <i>Proteus</i> species	Salm C1 - <i>Salmonella</i> species group C1
Ente - <i>Enterobacter</i> species	Prp - <i>Propionibacterium</i> species	Salm C2 - <i>Salmonella</i> species group C2
Erys - <i>Erysipelothrix</i>	Pseu - <i>Pseudomonas</i> species	Salm D - <i>Salmonella</i> species group D
Fus - <i>Fusobacterium</i>	R equ - <i>Rhodococcus equi</i>	Salm E - <i>Salmonella</i> species group E
G ana - <i>Gallibacterium anatis</i>		