

Data reported as: % susceptible (# isolates tested)¹

	B bron	E coli	E fael	E faem	Ente	K pneu	P aer	P mult	Pseu	S aur	S can	S pint
Amikacin	100% (10)	100% (60)	37% (27)	33% (6)	92% (13)	100% (2)	100% (24)	84% (25)	92% (12)	100% (16)	15% (13)	100% (13)
Amoxicillin/ Clavulanic Acid	100% (10)	75% (60)	93% (27)	33% (6)	31% (13)	100% (2)	0% (24)	100% (25)	40% (12)	100% (16)	100% (13)	92% (13)
Ampicillin	71% (10)	63% (60)	96% (27)	33% (6)	46% (13)	0% (2)	4% (24)	100% (25)	40% (12)	44% (16)	100% (13)	62% (13)
Cefazolin	0% (10)	88% (60)	4% (27)	17% (6)	23% (13)	100% (2)	0% (24)	100% (25)	25% (12)	100% (16)	100% (13)	92% (13)
Cefovecin	0% (9)	89% (46)	0% (24)	0% (5)	77% (13)	100% (2)	0% (23)	96% (24)	25% (12)	100% (13)	100% (11)	90% (10)
Cefoxitin	0% (10)	88% (60)	4% (27)	0% (6)	31% (13)	100% (2)	0% (24)	96% (25)	42% (12)	75% (16)	100% (13)	92% (13)
Cefpodoxime	0% (10)	85% (60)	15% (27)	0% (6)	77% (13)	100% (2)	0% (24)	96% (25)	25% (12)	94% (16)	100% (13)	92% (13)
Ceftiofur	0% (10)	87% (60)	4% (27)	0% (6)	77% (13)	100% (2)	4% (24)	100% (25)	33% (12)	100% (16)	100% (13)	92% (13)
Cephalothin	0% (4)	72% (25)	0% (11)	100% (1)	0% (1)	ND	0% (5)	100% (9)	0% (2)	100% (9)	100% (6)	86% (7)
Chloramphenicol	100% (10)	87% (60)	89% (27)	100% (6)	69% (13)	100% (2)	4% (24)	100% (25)	33% (12)	69% (16)	100% (13)	100% (13)
Clindamycin	0% (10)	0% (60)	7% (27)	17% (6)	0% (13)	0% (2)	0% (24)	0% (25)	0% (12)	100% (16)	77% (13)	69% (13)
Doxycycline	100% (9)	87% (46)	92% (24)	60% (5)	77% (13)	100% (2)	17% (23)	96% (24)	58% (12)	100% (13)	73% (11)	50% (10)
Enrofloxacin	80% (10)	85% (60)	37% (27)	0% (6)	92% (13)	100% (2)	79% (24)	100% (25)	92% (12)	81% (16)	46% (13)	62% (13)
Erythromycin	0% (10)	0% (60)	11% (27)	0% (6)	0% (13)	0% (2)	0% (24)	12% (25)	25% (12)	63% (16)	0% (13)	54% (13)
Gentamicin	100% (10)	97% (60)	67% (27)	17% (6)	92% (13)	100% (2)	92% (24)	96% (25)	92% (12)	100% (16)	69% (13)	85% (13)
Imipenem	100% (10)	100% (60)	96% (27)	33% (6)	92% (13)	100% (2)	100% (24)	100% (25)	92% (12)	100% (16)	100% (13)	92% (13)
Marbofloxacin	100% (10)	92% (60)	45% (27)	0% (6)	92% (13)	100% (2)	100% (24)	100% (25)	100% (12)	94% (16)	77% (13)	69% (13)
Orbifloxacin	100% (1)	86% (14)	0% (3)	0% (1)	ND	ND	0% (1)	100% (1)	ND	100% (3)	0% (2)	67% (3)
Oxacillin ³	NA	NA	NA	NA	NA	NA	NA	NA	NA	100% (16)	NA	92% (13)
Penicillin	0% (10)	0% (60)	96% (27)	33% (6)	0% (13)	0% (2)	0% (24)	54% (25)	0% (12)	38% (16)	100% (13)	38% (13)
Tetracycline	100% (1)	79% (14)	33% (3)	100% (1)	ND	ND	0% (1)	100% (1)	ND	100% (3)	50% (2)	100% (3)
Ticarcillin	70% (10)	72% (60)	11% (27)	17% (6)	69% (13)	0% (2)	96% (24)	100% (25)	75% (12)	100% (16)	100% (13)	92% (13)
Ticarcillin/ Clavulanic Acid	100% (10)	87% (60)	11% (27)	17% (6)	77% (13)	100% (2)	96% (24)	100% (25)	83% (12)	100% (16)	100% (13)	92% (13)
Trimethoprim/ Sulphamethoxazole	60% (10)	90% (60)	93% (27)	100% (6)	92% (13)	100% (2)	21% (24)	92% (25)	58% (12)	100% (16)	100% (13)	77% (13)

³ Isolates resistant to oxacillin are interpreted as methicillin resistant.

Key:

1	Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility	
2	See Salmonella serotype table for most common serotypes isolated within each group	
3	Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.	
4	A result of ≤ 2 ug/ml for Carbadox is a conservative indicator of bacterial inhibition by this antimicrobial agent. The result shown is based on pharmacokinetic research indicating an average Carbadox level of 4.5 mcg/ml in the small intestine of pigs fed a dose rate of 50 g/ton. (De Graff 1988).	
5	Multidrug resistant isolates were found resistant to most classes of antimicrobial in the 1 st round of testing. This table represents additional Disk Diffusion testing for those isolates.	
NA	Not applicable	
ND	Not done	
NI	No interpretation	
A equ - Actinobacillus equuli	H ecol - hemolytic E. coli	S aur - Staphylococcus aureus
A suis - Actinobacillus suis	H som - Histophilus somni	S beta- Beta Streptococcus species
Abua - Acinetobacter species	HPS - Haemophilus parasuis	S can - Streptococcus canis
Amy - Actinomyces species	K pneu - Klebsiella pneumoniae	S chol - Salmonella choleraesuis
APP - Actinobacillus pleuropneumoniae	M bov - Moraxella bovis	S dysg - Streptococcus dysgalactiae
B bron - Bordetella bronchiseptica	M haem - Mannheimia haemolytica	S epi- Staphylococcus epidermidis
B tre - Bibersteinia trehalosi (formerly Pasteurella trehalosi)	P aer - Pseudomonas aeruginosa	S equi - Streptococcus equi
Bact - Bacteroides group	P cab - Pasteurella caballi	S equus - Streptococcus equisimilis
C diff - Clostridium difficile	P mult - Pasteurella multocida	S pint - Staph pseudintermedius
C perf - Clostridium perfringens	Past - Pasteurella species	S suis - Streptococcus suis
Clos - Clostridium species	Pec - Peptococcus species	S ube - Streptococcus uberis
E coli - Escherichia coli	Pes - Peptostreptococcus species	S zoo - Streptococcus zooepidemicus
E fael - Enterococcus faecalis	Pmul A - Pasteurella multocida Type A	Salm sp- Salmonella species
E faem - Enterococcus faecium	Pmul D - Pasteurella multocida Type D	Salm B - Salmonella species group B
Enc - Enterococcus species	Prot - Proteus species	Salm C1 - Salmonella species group C1
Ente - Enterobacter species	Prp - Propionibacterium species	Salm C2 - Salmonella species group C2
Erys - Erysipelothrix	Pseu - Pseudomonas species	Salm D - Salmonella species group D
Fus - Fusobacterium	R equ - Rhodococcus equi	Salm E - Salmonella species group E
G ana - Gallibacterium anatis		