

*Data reported as: % susceptible (# isolates tested)<sup>1</sup>*

Antibiotic	E coli	E faem	Ente	K pneu	P mult	S aur	S dysg	S epi	S ube
Ampicillin	80% (49)	100% (2)	63% (8)	0% (10)	100% (2)	78% (64)	100% (27)	83% (36)	78% (40)
Ceftiofur	96% (49)	50% (2)	88% (8)	100% (10)	100% (2)	100% (64)	100% (27)	94% (36)	85% (40)
Cephalothin	82% (49)	50% (2)	0% (8)	100% (10)	100% (2)	100% (64)	100% (27)	94% (36)	100% (40)
Erythromycin	0% (49)	0% (2)	0% (8)	0% (10)	0% (2)	97% (64)	93% (27)	83% (36)	75% (40)
Oxacillin	0% (49)	0% (2)	0% (8)	0% (10)	50% (2)	94% (64)	96% (27)	6% (36)	78% (40)
Penicillin	0% (49)	50% (2)	0% (8)	0% (10)	100% (2)	77% (64)	100% (27)	81% (36)	35% (40)
Penicillin/Novobiocin	0% (49)	NI	NI	0% (10)	NI	97% (64)	100% (27)	94% (36)	98% (40)
Pirlimycin	0% (49)	50% (2)	0% (8)	0% (10)	0% (2)	97% (64)	93% (27)	86% (36)	78% (40)
Sulfadimethoxine	43% (49)	0% (2)	0% (8)	30% (10)	0% (2)	88% (64)	59% (27)	69% (36)	0% (40)
Tetracycline	47% (49)	50% (2)	50% (8)	80% (10)	50% (2)	89% (64)	19% (27)	61% (36)	65% (40)

<sup>3</sup> Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.