

Data reported as: % susceptible (# isolates tested)¹

Antibiotic	B bron	E coli	E fael	E faem	Ente	K pneu	P aer	P mult	Pseu	S aur	S can	S pint
Amikacin	100% (19)	98% (684)	22% (225)	7% (44)	100% (112)	96% (52)	93% (248)	100% (36)	89% (105)	100% (34)	3% (233)	99% (812)
Amoxicillin/Clavulanic Acid	74% (19)	77% (684)	97% (225)	36% (44)	35% (112)	65% (52)	2% (248)	100% (36)	46% (105)	76% (34)	100% (233)	86% (812)
Ampicillin	16% (19)	62% (684)	96% (225)	34% (44)	40% (112)	13% (52)	1% (248)	100% (36)	43% (105)	26% (34)	100% (233)	48% (812)
Cefazolin	5% (19)	84% (684)	4% (225)	5% (44)	24% (112)	63% (52)	2% (248)	100% (36)	36% (105)	76% (34)	100% (233)	86% (812)
Cefovecin	5% (19)	84% (684)	3% (225)	2% (44)	79% (112)	79% (52)	1% (248)	100% (36)	41% (105)	76% (34)	98% (233)	82% (812)
Cefoxitin	5% (19)	86% (684)	1% (225)	0% (44)	40% (112)	67% (52)	2% (248)	100% (36)	39% (105)	53% (34)	100% (233)	84% (812)
Cefpodoxime	5% (19)	84% (684)	16% (225)	5% (44)	81% (112)	81% (52)	1% (248)	97% (36)	39% (105)	68% (34)	98% (233)	83% (812)
Ceftiofur	0% (19)	84% (684)	7% (225)	5% (44)	83% (112)	81% (52)	1% (248)	100% (36)	41% (105)	76% (34)	99% (233)	84% (812)
Cephalothin	83% (6)	64% (53)	7% (98)	18% (17)	10% (10)	0% (2)	0% (22)	100% (2)	27% (11)	75% (28)	99% (201)	86% (664)
Chloramphenicol	95% (19)	84% (684)	92% (225)	93% (44)	79% (112)	83% (52)	2% (248)	100% (36)	46% (105)	76% (34)	97% (233)	87% (812)
Clindamycin	5% (19)	0% (684)	1% (225)	32% (44)	0% (112)	0% (52)	0% (248)	0% (36)	10% (105)	88% (34)	88% (233)	74% (812)
Doxycycline	100% (19)	82% (684)	73% (225)	45% (44)	71% (112)	77% (52)	5% (248)	97% (36)	70% (105)	85% (34)	64% (233)	61% (812)
Enrofloxacin	95% (19)	88% (684)	32% (225)	0% (44)	96% (112)	85% (52)	43% (248)	100% (36)	62% (105)	82% (34)	48% (233)	78% (812)
Erythromycin	0% (19)	0% (684)	28% (225)	5% (44)	1% (112)	0% (52)	0% (248)	14% (36)	25% (105)	65% (34)	0% (233)	73% (812)
Gentamicin	79% (19)	91% (684)	47% (225)	2% (44)	97% (112)	90% (52)	75% (248)	100% (36)	86% (105)	97% (34)	42% (233)	81% (812)
Imipenem	100% (19)	100% (684)	99% (225)	32% (44)	100% (112)	100% (52)	97% (248)	100% (36)	95% (105)	74% (34)	100% (233)	86% (812)
Marbofloxacin	95% (19)	90% (684)	37% (225)	0% (44)	96% (112)	94% (52)	74% (248)	100% (36)	88% (105)	82% (34)	82% (233)	83% (812)
Oxacillin ³	NI	NI	NI	NI	NI	NI	NI	NI	NI	76% (34)	NI	82% (812)
Penicillin	0% (19)	0% (684)	97% (225)	32% (44)	0% (112)	0% (52)	0% (248)	81% (36)	1% (105)	21% (34)	98% (233)	31% (812)
Ticarcillin	47% (19)	67% (684)	8% (225)	9% (44)	77% (112)	19% (52)	90% (248)	100% (36)	71% (105)	74% (34)	100% (233)	85% (812)
Ticarcillin/Clavulanic Acid	95% (19)	80% (684)	8% (225)	7% (44)	88% (112)	81% (52)	91% (248)	100% (36)	77% (105)	76% (34)	99% (233)	85% (812)
Trimethoprim/Sulphamethoxazole	37% (19)	87% (684)	90% (225)	82% (44)	97% (112)	79% (52)	7% (248)	97% (36)	50% (105)	100% (34)	97% (233)	78% (812)

³ Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.

Key:

1	Data is reported as: % susceptible (# isolates tested) - not all bacteria isolated at ISU VDL have been tested for antimicrobial susceptibility	
2	See Salmonella serotype table for most common serotypes isolated within each group	
3	Isolates resistant to oxacillin are interpreted as potentially methicillin resistant.	
4	A result of ≤ 2 ug/ml for Carbadox is a conservative indicator of bacterial inhibition by this antimicrobial agent. The result shown is based on pharmacokinetic research indicating an average Carbadox level of 4.5 mcg/ml in the small intestine of pigs fed a dose rate of 50 g/ton. (De Graff 1988).	
5	Multidrug resistant isolates were found resistant to most classes of antimicrobial in the 1 st round of testing. This table represents additional Disk Diffusion testing for those isolates.	
NA	Not applicable	
ND	Not done	
NI	No interpretation	
A equ - Actinobacillus equuli	H ecol - hemolytic E.coli	S aur - Staphylococcus aureus
A suis - Actinobacillus suis	H som - Histophilus somni	S beta- Beta Streptococcus species
Abua - Acinetobacter species	HPS - Haemophilus parasuis	S can - Streptococcus canis
Amy - Actinomyces species	K pneu - Klebsiella pneumoniae	S chol - Salmonella choleraesuis
APP - Actinobacillus pleuropneumoniae	M bov - Moraxella bovis	S dysg - Streptococcus dysgalactiae
B bron - Bordetella bronchiseptica	M haem - Mannheimia haemolytica	S epi- Staphylococcus epidermidis
B tre - Bibersteinia trehalosi (formerly Pasteurella trehalosi)	P aer - Pseudomonas aeruginosa	S equi - Streptococcus equi
Bact - Bacteroides group	P cab - Pasteurella caballi	S equus - Streptococcus equisimilis
C diff - Clostridium difficile	P mult - Pasteurella multocida	S pint - Staph pseudintermedius
C perf - Clostridium perfringens	Past - Pasteurella species	S suis - Streptococcus suis
Clos - Clostridium species	Pec - Peptococcus species	S ube - Streptococcus uberis
E coli - Escherichia coli	Pes - Peptostreptococcus species	S zoo - Streptococcus zooepidemicus
E fael - Enterococcus faecalis	Pmul A - Pasteurella multocida Type A	Salm sp- Salmonella species
E faem - Enterococcus faecium	Pmul D - Pasteurella multocida Type D	Salm B - Salmonella species group B
Enc - Enterococcus species	Prot - Proteus species	Salm C1 - Salmonella species group C1
Ente - Enterobacter species	Prp - Propionibacterium species	Salm C2 - Salmonella species group C2
Erys - Erysipelothrix	Pseu - Pseudomonas species	Salm D - Salmonella species group D
Fus - Fusobacterium	R equ - Rhodococcus equi	Salm E - Salmonella species group E
G ana - Gallibacterium anatis		